

# Prenatal Infection and Neurodevelopmental Disabilities

**Prenatal infection** and mediators of inflammation are risk factors for **neurodevelopmental** disabilities, such as **cerebral palsy** and **autism**.

General Information	
<b>Broad Focus Area</b>	Neurodevelopment and behavior
<b>Background and Justification</b>	Cerebral palsy and autism are uncommon but serious developmental disabilities that have a dramatic effect on the lives of the affected persons and their families. Exposure to prenatal infection or mediators of inflammation may increase risk of cerebral palsy. <sup>1</sup> Fetal inflammatory response to chorioamnionitis (intrauterine infection) includes increased levels of fetal cytokines; such cytokines can be neurotoxic. <sup>2,3</sup> While few studies of chorioamnionitis and cerebral palsy among children born at term have been done, <sup>4,5</sup> it has been estimated that about 28% of cerebral palsy in preterm infants and 12% of cerebral palsy in term infants may be due to chorioamnionitis. <sup>6</sup> A recent study by Wu et al. observed a four-fold increased risk (adjusted) of cerebral palsy in subjects with diagnosed chorioamnionitis. <sup>7</sup> While there are some data on the relation of viral infections in pregnancy to occurrence of autism, few causal agents have been established—rubella being one. <sup>8</sup> The more general relation of prenatal infection, such as chorioamnionitis, and of exposure to mediators of inflammation, to risk of autism has not been studied. Such studies are overdue because of the role immune abnormalities may play in autism and the increased knowledge of the neurotoxicity of inflammatory cytokines. <sup>9</sup> Although a large portion of autism may be genetically determined, the inherited predisposition may increase susceptibility to infection or inflammatory-induced disease.
<b>Prevalence/ Incidence</b>	Cerebral palsy affects approximately 0.2% of children, <sup>10</sup> and autism affects about 0.3%. <sup>11</sup> Whether the frequency of autism is increasing is controversial, because recent estimates of higher prevalence may be due to inclusion of less severe cases. In term pregnancies, about 1-2 percent are affected by chorioamnionitis; in pregnancies ending in preterm births, the prevalence of such infection is higher. <sup>6</sup>
<b>Economic Impact</b>	While no studies have precisely calculated all of the costs associated with autism, a U.K. report estimates the lifetime custodial costs of autism spectrum disorders in the range of \$3-\$4 million per child, with societal costs likely to be triple the individual estimate. <sup>12, 13</sup> The lifetime economic costs of cerebral palsy have been estimated at \$11.5 billion per annual cohort. <sup>14</sup>

Exposure Measures		Outcome Measures	
<b>Primary/ Maternal</b>	Maternal infection/inflammation: - Infection serology (lymphocytes, antibodies, cytokines/interleukins, inflammatory markers) - Medical history of fever and infection (medicine usage) - Dental exams	<b>Primary/ Child</b>	Neurological development
Methods	- Blood samples - Vaginal/cervical cultures - Examination by a medical	Methods	Direct observation by medical professional: fetal ultrasound, neurological exam, autism

	professional - Interview			screening test
Life Stage	Repeated measures 1 <sup>st</sup> through 3 <sup>rd</sup> trimesters and birth		Life Stage	Prenatal through year 7
<b>Primary/ Child</b>	Prenatal infection/inflammation: - Infection serology (lymphocytes, antibodies, cytokines/interleukins, inflammatory markers) - Umbilical cord/placental (antibodies; cytokines)		<b>Secondary/ Child</b>	School performance
Methods	- Umbilical cord blood culture/pathology		Methods	School record examination for grades/performance
Life Stage	Birth		Life Stage	Follow-up in year 7
<b>Secondary/ Maternal</b>	Retrospective Medical Record Review			
Methods	Medical and obstetrical history, family history			
Life Stage	Repeated measures 1 <sup>st</sup> through 3 <sup>rd</sup> trimesters and birth			

Important Confounders/Covariates	
Family history	Twin and family studies have suggested a genetic link, which may be shown through family histories, to these disorders. <sup>15</sup>
Mother's medical and obstetrical histories	Particularly in cases without a family history, various obstetric complications are often cited as possible causes for fetal neurodevelopmental disruption and consequent disorders. <sup>1</sup>

Population of Interest	Estimated Effect that is Detectable
All children	Assuming 100,000 infants born into the study, with an exposure prevalence of 2%, the smallest detectable relative risk would be, for cerebral palsy, 2.8; and for autism, 2.4.

Other Design Issues	
Cost/Complexity of Data Collection	Retention of index children at least to or beyond the average age of diagnosis for these disorders (e.g., age 6 or 7) will be important to address this hypothesis with sufficient power. Necessity of umbilical cord blood samples or samples taken early in infancy may have a very strong impact as there would need for coordination with medical professionals.
Cost of Sample Analysis	Because the contacts with these patients will essentially fall within the scope of standard of care during the pregnancy, additional cost will be relatively minor and entail simply maintaining the pregnancy and infant/childhood data-base and the tissue sample repositories.

## References:

- <sup>1</sup> Nelson KB, Willoughby RE. Overview: infection during pregnancy and neurologic outcome in the child. *Ment Retard Dev Disabil Res Rev.* 2002;8(1):1-2.
- <sup>2</sup> Yoon BH, Romero R, Park JS, Chang JW, Kim YA, Kim JC, Kim KS. Microbial invasion of the amniotic cavity with *Ureaplasma urealyticum* is associated with a robust host response in fetal, amniotic, and maternal compartments. *Am J Obstet Gynecol.* 1998 Nov;179(5):1254-60.
- <sup>3</sup> Dammann O, Leviton A. Infection remote from the brain, neonatal white matter damage, and cerebral palsy in the preterm infant. *Semin Pediatr Neurol.* 1998 Sep; 5(3):190-201.
- <sup>4</sup> Grether JK, Nelson KB. Maternal infection and cerebral palsy in infants of normal birth weight. *JAMA.* 1997 Jul 16;278(3):207-11.
- <sup>5</sup> Nelson KB, Ellenberg JH. Antecedents of cerebral palsy. I. Univariate analysis of risks. *Am J Dis Child.* 1985 Oct;139(10):1031-8.
- <sup>6</sup> Wu YW, Colford JM Jr. Chorioamnionitis as a risk factor for cerebral palsy: A meta-analysis. *JAMA.* 2000 Sep 20;284(11):1417-24.
- <sup>7</sup> Wu, Y.W., Escobar, G.J., Grether, J.K., Croen, L.A., Greene, J.D., and T.B. Newman. 2003. Chorioamnionitis and cerebral palsy in term and near-term infants. *JAMA* 290(20): 2677-84.
- <sup>8</sup> Rodier PM, Hyman SL. Early environmental factors in autism. *MENT RETARD DEV D R* 4 (2): 121-128 1998.
- <sup>9</sup> Horning M, Lipkin WI. *MRDD Research Reviews* 2001;200-210.
- <sup>10</sup> Kuban KC, Leviton A. Cerebral palsy. *N Engl J Med.* 1994 Jan 20;330(3):188-95.
- <sup>11</sup> Yeargin-Allsopp M, Rice C, Karapurkar T, Doernberg N, Boyle C, Murphy C. Prevalence of autism in a US metropolitan area. *JAMA.* 2003 Jan 1;289(1):49-55.
- <sup>12</sup> International Child Development Resource Center (ICDRC). ICDRC perspective of the epidemic of autism spectrum disorders: a view from the trenches. Executive Briefing of Claude A. Allen, Deputy Secretary of Health and Human Services, June 4, 2002. Available online: [http://www.icdrc.org/executive\\_briefing.html](http://www.icdrc.org/executive_briefing.html).
- <sup>13</sup> Jarbrink, K. and M. Knapp. The economic impact of autism in Britain. *Autism* 2001 Mar; 5(1):7-22.
- <sup>14</sup> MMWR. Economic Costs Associated with Mental Retardation, Cerebral Palsy, Hearing Loss, and Vision Impairment --- United States, 2003. January 30, 2004; 53(03): 57-59.
- <sup>15</sup> Muhle R., Trentacoste S.V, Rapin I. The Genetics of Autism. *PEDIATRICS* 113(5) May 2004, pp. e472-e486.